

COVID-19 Vaccine Eligibility

Use this chart to help determine when to give COVID-19 vaccine based on age or health conditions.

Dose	Age:	5-11	12-17	18+
Pfizer–Pediatric (5-11)	Primary 2 doses (21 days apart)	✓		
	Additional (3rd) dose for immunocompromised , at least 28 days after 2nd Pfizer dose	✓		
Pfizer/Comirnaty (12+)	Primary 2 doses (21 days apart)		✓	✓
	Additional (3rd) dose for immunocompromised , at least 28 days after 2nd Pfizer dose		✓	✓
	Booster dose* of Pfizer (or Moderna if age 18+), at least 5 months after 2nd dose of Pfizer		✓	✓
Moderna (18+)	Primary 2 doses (28 days apart)			✓
	Additional (3rd) dose for immunocompromised , at least 28 days after 2nd Moderna dose			✓
	Booster dose* of Moderna (half-dose) or Pfizer, at least 6 months after 2nd dose of Moderna			✓
Johnson & Johnson (18+)	Primary 1 dose			✓
	Booster dose* of different brand (preferred) or J&J, at least 2 months after primary J&J dose			✓
Use of Pfizer or Moderna is preferred.*				✓

* Use of mRNA COVID-19 vaccines is preferred over the Janssen COVID-19 vaccine. However, the Janssen vaccine may be offered in some situations:

- When there is a contraindication to mRNA COVID-19 vaccines (e.g., severe allergic reaction after a previous dose or to a component of an mRNA COVID-19 vaccine)
- When a person would otherwise remain unvaccinated due to limited access to mRNA COVID-19 vaccines
- When a person wants to receive the Janssen COVID-19 Vaccine despite the safety concerns identified

View [Related Guidance and Requirements](#). (Eligibility is subject to change as guidance is updated.)